Protect yourself away from home

- Avoid social and community gatherings where 10 people or more would come into close contact.
- Practice <u>social distancing</u> advice below when you are in a group of people.
- Avoid unnecessary (non-urgent) air, bus or train travel.
- Limit visitation to older relatives or friends (especially in nursing or care homes).

Reported illnesses have ranged from mild symptoms to severe illness and death for confirmed coronavirus disease 2019 (COVID-19) cases.

These symptoms may appear **2-14 days after exposure** (based on the incubation period of MERS-CoV viruses).

- Fever
- Cough
- Shortness of breath

If you develop **emergency warning signs** for COVID-19 get **medical attention immediately**. Emergency warning signs include*:

- Trouble breathing
- Persistent pain or pressure in the chest
- New confusion or inability to arouse
- Bluish lips or face

*This list is not all inclusive. Please consult your medical provider for any other symptoms that are severe or concerning.

Know How it Spreads



- There is currently no vaccine to prevent coronavirus disease 2019 (COVID-19).
- The best way to prevent illness is to avoid being exposed to this virus.
- The virus is thought to spread mainly from person-to-person.
 - Between people who are in close contact with one another (within about 6 feet).
 - Through respiratory droplets produced when an infected person coughs or sneezes.
- These droplets can land in the mouths or noses of people who are nearby or possibly be inhaled into the lungs.

Take steps to protect yourself



Clean your hands often

- **Wash your hands** often with soap and water for at least 20 seconds especially after you have been in a public place, or after blowing your nose, coughing, or sneezing.
- If soap and water are not readily available, use a hand sanitizer that contains at least 60% alcohol. Cover all surfaces of your hands and rub them together until they feel dry.
- Avoid touching your eyes, nose, and mouth with unwashed hands.



Avoid close contact

- Avoid close contact with people who are sick
- Put **distance between yourself and other people** if COVID-19 is spreading in your community. This is especially important for <u>people who are at higher risk of getting very sick</u>.

Take steps to protect others



Stay home if you're sick

• **Stay home,** if you are sick, call your primary care physician or the ER to minimize exposure of yourself and others.



Cover coughs and sneezes

- **Cover your mouth and nose** with a tissue when you cough or sneeze or use the inside of your elbow.
- Throw used tissues in the trash.
- Immediately **wash your hands** with soap and water for at least 20 seconds. If soap and water are not readily available, clean your hands with a hand sanitizer that contains at least 60% alcohol.



Wear a facemask if you are sick

• If you are sick: You should wear a facemask when you are around other people (e.g., sharing a room or vehicle) and before you enter a healthcare provider's office. If you are not able to wear a facemask (for example, because it causes trouble breathing), then you should do your best to cover your coughs and sneezes, and people who are caring for you should wear a facemask if they enter your room. Learn what to do if you are sick.

• If you are NOT sick: You do not need to wear a facemask unless you are caring for someone who is sick (and they are not able to wear a facemask). Facemasks may be in short supply and they should be saved for caregivers.



Clean and disinfect

- Clean AND disinfect <u>frequently touched surfaces</u> daily. This includes tables, doorknobs, light switches, countertops, handles, desks, phones, keyboards, toilets, faucets, and sinks.
- If surfaces are dirty, clean them: Use detergent or soap and water prior to disinfection.

To disinfect:

Most common EPA-registered household disinfectants will work. Use disinfectants appropriate for the surface.

Options include:

• Diluting your household bleach.

To make a bleach solution, mix:

- 5 tablespoons (1/3rd cup) bleach per gallon of water OR
- o 4 teaspoons bleach per quart of water

Follow manufacturer's instructions for application and proper ventilation. Check to ensure the product is not past its expiration date. Never mix household bleach with ammonia or any other cleanser. Unexpired household bleach will be effective against coronaviruses when properly diluted.

Alcohol solutions.

Ensure solution has at least 70% alcohol.

• Other common EPA-registered household disinfectants.

Products with <u>EPA-approved emerging viral pathogens pdf icon</u>[7 pages]<u>external icon</u> claims are expected to be effective against COVID-19 based on data for harder to kill viruses. Follow the manufacturer's instructions for all cleaning and disinfection products (e.g., concentration, application method and contact time, etc.).